



GEOLOGY OF THE INTERMOUNTAIN WEST

an open-access journal of the Utah Geological Association

ISSN 2380-7601

Volume 11

2024

THE FIRST DINOSAUR POSTCRANIAL BODY FOSSILS FROM THE LOWER JURASSIC KAYENTA FORMATION OF UTAH

Adam D. Marsh, Donald D. De Blieux, and James I. Kirkland



Illustration by Brian Engh
LivingRelicProductions.com

This is an open-access article in which the Utah Geological Association permits unrestricted use, distribution, and reproduction of text and figures that are not noted as copyrighted, provided the original author and source are credited. Email inquiries to GIW@utahgeology.org.



GEOLOGY OF THE INTERMOUNTAIN WEST

an open-access journal of the Utah Geological Association

ISSN 2380-7601

Volume 11

2024

Editors

Douglas A. Sprinkel Azteca Geosolutions 801.391.1977 GIW@utahgeology.org dsprinkel@gmail.com	Steven Schamel GeoX Consulting, Inc. 801.583.1146 geox-slc@comcast.net
Thomas C. Chidsey, Jr. Utah Geological Survey, Emeritus 801.824.0738 tomchidsey@gmail.com	John R. Foster Utah Field House of Natural History State Park Museum 435.789.3799 johnfoster@utah.gov
Bart J. Kowallis Brigham Young University 801.380.2736 bkowallis@gmail.com	William R. Lund Utah Geological Survey, Emeritus 435.590.1338 williamlundugs@gmail.com

Production

Cover Design and Desktop Publishing
Douglas A. Sprinkel

Cover

A theropod dinosaur scans the horizon as sauropod-morphs walk to the river, which flows across the Early Jurassic landscape. The sediments deposited by the river systems will eventually become the Kayenta Formation. Artwork by Brian Eng, livingrelicproductions.com used by permission.



Geology of the Intermountain West (GIW) is an open-access journal in which the Utah Geological Association permits unrestricted use, distribution, and reproduction of text and figures that are not noted as copyrighted, provided the original author and source are credited.

2023–2024 UGA Board

President	Eugene Syzanski	eugenes@utah.gov	801.537.3364
President-Elect	Keilee Higgs	keileeann@utah.gov	801.537.3304
Program Chair	Chris Stallard	cstallard@utah.gov	801.386.0976
Treasurer	Aubry DeReuil	aubry@zanskar.us	850.572.2543
Secretary	Trae Boman	tboman@teanues.com	801.648.5206
Past President	Rick Ford	rford@weber.edu	801.915.3188

UGA Committees

Environmental Affairs	Craig Eaton	eaton@ihi-env.com	801.633.9396
Geologic Road Sign	Greg Gavin	greg@loughlinwater.com	801.541.6258
Historian	Paul Anderson	paul@pbageo.com	801.364.6613
Outreach	Greg Nielsen	gnielsen@weber.edu	801.626.6394
Public Education	Zach Anderson	zanderson@utah.gov	801.537.3300
	Matt Affolter	gfl247@yahoo.com	
Publications	Paul Inkenbrandt	paulinkenbrandt@utah.gov	801.537.3361
Publicity	Paul Inkenbrandt	paulinkenbrandt@utah.gov	801.537.3361
Social/Recreation	Roger Bon	rogerbon@xmission.com	801.942.0533

AAPG House of Delegates

2023–2026 Term	David A. Wavrek	dwavrek@petroleumsystems.com	801.322.2915
----------------	-----------------	------------------------------	--------------

State Mapping Advisory Committee

UGA Representative	Bill Loughlin	bill@loughlinwater.com	435.649.4005
--------------------	---------------	------------------------	--------------

Earthquake Safety Committee

Chair	Grant Willis	gwillisgeol@gmail.com	801.537.3355
-------	--------------	-----------------------	--------------

UGA Website — www.utahgeology.org

Webmaster	Paul Inkenbrandt	paulinkenbrandt@utah.gov	801.537.3361
-----------	------------------	--------------------------	--------------

Scholarship Golf Tournament

Co-Chair	Rick Ford	rford@weber.edu	801.915.3188
Co-Chair	John South	jsouth@utah.gov	385.266.2113

UGA Newsletter

Newsletter Editor	William Lund	uga.newsletter@gmail.com	435.590.1338
-------------------	--------------	--------------------------	--------------

Become a member of the UGA to help support the work of the Association and receive notices for monthly meetings, annual field conferences, and new publications. Annual membership is \$30 and annual student membership is only \$5. Visit the UGA website at www.utahgeology.org for information and membership application.

The UGA board is elected annually by a voting process through UGA members. However, the UGA is a volunteer-driven organization, and we welcome your voluntary service. If you would like to participate please contact the current president or committee member corresponding with the area in which you would like to volunteer.



The First Dinosaur Postcranial Body Fossils from the Lower Jurassic Kayenta Formation of Utah

Adam D. Marsh¹, Donald D. De Blieux², and James I. Kirkland²

¹Department of Science and Resource Management, Petrified Forest National Park, AZ 86028 USA; adam_marshall@nps.gov

²Utah Geological Survey, Salt Lake City, UT 84114 USA; dondeblieux@utah.gov, jameskirkland@utah.gov

ABSTRACT

The vertebrate assemblage of the Lower Jurassic Kayenta Formation is known for its preservation of post-end Triassic mass extinction lineages, including lissamphibians, lepidosaurs, turtles, mammali-amorphs, crocodylomorphs, pterosaurs, and ornithischian, theropod, and sauropodomorph dinosaurs. Most of the body fossils from the formation are known from its ‘silty facies’ in north-central Arizona and southwestern Utah, whereas the sandier ‘typical facies’ of northeastern Arizona preserves few body fossils, and until recently they were completely absent in the typical facies of southeastern Utah. A 2011 team conducting a paleontological survey of Arches National Park discovered the first body fossils from the typical facies of the Kayenta Formation in Utah, here identified as belonging to a single individual of a saurischian dinosaur, likely a theropod. The fossil elements include a partial centrum articular face, a prezygapophysis, part of a caudal vertebra, the distal end of a left radius, part of the distal end of a left femur, a shaft fragment from the left fibula, the distal end of right metatarsal I, and the proximal portion of left metatarsals III and IV. This specimen from Arches National Park underscores the importance of federally protected land in fossil resource management and suggests that the typical facies of the Kayenta Formation may be under-sampled and could preserve more vertebrate bones than previously thought.

INTRODUCTION

The Lower Jurassic Kayenta Formation unconformably overlies the Upper Triassic-Lower Jurassic Moenave Formation or Wingate Sandstone, depending on the location (the J-sub-K unconformity; Riggs and Blakey, 1993; Blakey, 1994). It interfingers with the higher Navajo Sandstone (Figure 1A), forming a sequence of strata that records erg sands blowing over fluviolacustrine deposits as the North American plate moved north into an arid climate belt (Harshbarger et al., 1957; Riggs et al., 1993; Blakey, 1994; Priddy and Clarke, 2020). The Kayenta Formation preserves an important vertebrate assemblage that

represents the terrestrial biotic recovery following the end-Triassic mass extinction in Western North America. Fossils discovered in the formation include lissamphibians, lepidosaurs, turtles, mammaliomorphs, pterosaurs, crocodylomorphs, and dinosaurs (Clark and Fastovsky, 1986; Sues et al., 1994; Lucas et al., 2005; Tykoski, 2005). This assemblage is best known from the ‘silty facies’ of parts of northern Arizona and southwestern Utah (Figure 1B), but its sandier ‘typical facies’ like what is present at its type section at Comb Ridge, Arizona, also preserves vertebrate fossils, primarily as tracks (Lockley and Hunt, 1994, 1995; Hunt and Lucas, 2006). As Harshbarger et al. (1957, p. 17) observed, “the typical and silty facies of the Kayenta

Citation for this article.

Marsh, A.D., De Blieux, D.D., and Kirkland, J.I., 2024, *The first dinosaur postcranial body fossils from the Lower Jurassic Kayenta Formation of Utah: Geology of the Intermountain West*, v. 11, p. 45–57, <https://doi.org/10.31711/giw.v11.pp45-57>.




Figure 1. Selected fossil localities in the Kayenta Formation of Arizona and Utah. (A) Map of northern Arizona and southern Utah showing localities and Arches National Park, the extent of Kayenta Formation facies is shown as brown and tan; (B) Stratigraphic context of the Glen Canyon Group near Arches National Park. Locality specimen ARCH 71v shown as a green star. Abbreviations: ARCH, Arches National Park; CR, Comb Ridge; GB, Goblin Valley; MF, Many Farms; sf, silty facies; TC, Tsegi Canyon; tf, typical facies; W, Warner Valley; WT, Ward Terrace, including the Echo and Adeii Eechi Cliffs; Z, Zion National Park. Approximate extent of the silty facies and typical facies of the Kayenta Formation modified from Priddy and Clarke (2020).

Formation represent a lateral variation of deposition within a single basin.”

This perceived disparity of the presence of body fossils in the two facies of the Kayenta Formation may be the result of different depositional processes, the fact that the typical facies tends to form cliffs rather than more accessible slopes, or geographic sampling bias. If the latter is true, then increased inventory of the typical facies of the Kayenta Formation should result in the discovery of more body fossils. This hypothesis was tested by a paleontological inventory that was undertaken within Arches National Park (ARCH) in 2011, resulting in the discovery of fragmentary fossil bones from the Kayenta Formation (Figure 2). Here, we describe that specimen (ARCH 4012) as the first dinosaur bones from the typical facies of the Kayenta Formation in Utah. This specimen highlights the importance of fossil resources on federally protected land and suggests that more vertebrate bones may be present elsewhere in the typical facies of the Kayenta Formation.

MATERIALS AND METHODS

A two-week paleontological field survey of the Phanerozoic strata of ARCH was led by the Utah Geological Survey (UGS) in 2011 under cooperative agreement #H230097080 with the U.S. National Park Service (NPS; Geologic Resources Division). A previous NPS ARCH survey was accomplished in 2000 (Swanson et al., 2005) but it focused on the Chinle Formation (Upper Triassic), Curtis Formation (Middle Jurassic), Morrison Formation (Upper Jurassic), and Cedar Mountain Formation (Lower Cretaceous). Paleontological localities identified or monitored during the 2011 survey were photographed and documented in the NPS and UGS databases (Madsen et al., 2012), and scientifically significant fossils were collected, including ARCH 4012. The ARCH collection, formerly at a regional NPS facility in Moab, Utah, was rehoused with archival materials in 2020 and is repositated at Petrified Forest National Park (PEFO) with other fossil collections from the NPS

Southeastern Utah Group. Photographs of ARCH 4012 were taken with a Nikon D90 digital camera equipped with a Nikon AF-S NIKKOR 24–85 mm lens. ARCH 4012 was laser scanned at PEFO using an Artec Space Spider on Artec Studio 16 Professional (16.0.5.114). The resulting surface files can be accessed at <https://www.morphosource.org/projects/000572397>.

INSTITUTIONAL ABBREVIATIONS

ARCH, Arches National Park, Utah, USA.; MNA, Museum of Northern Arizona, Flagstaff, Arizona, USA.; PEFO, Petrified Forest National Park, Arizona, USA.; TMM, Texas Vertebrate Paleontology Collections, The University of Texas at Austin, Austin, Texas, USA.; UCMP, University of California Museum of Paleontology, Berkeley, California, USA.

SYSTEMATIC PALEONTOLOGY

AVEMETATARSALIA Benton, 1999 sensu
Nesbitt et al., 2017

DINOSAURIA Owen, 1842 sensu
Padian and May, 1993

SAURISCHIA Seeley, 1887 sensu
Gauthier, 1986

THEROPODA Marsh, 1881
sensu Gauthier, 1986

(Figures 2 and 3)

Referred Specimen

Specimen ARCH 4012, fragmentary skeletal elements likely belonging to a single individual (Figures 2 and 3), including part of a presacral centrum and prezygapophysis, a caudal centrum, the distal end of a left radius, the distal end of a left femur, a shaft fragment from the proximal half of a left fibula, the distal end of right metatarsal I, and the proximal ends of left metatarsals III and IV.

Justification for Identification

The specimen can be attributed to the Saurischia

because of the presence of a hypantrum on the presacral vertebra, as well as the presence of a groove separating the lateral condyle and tibiofibular crest on the distal end of the femur (Figures 2P and 2R), both of which are present in early-diverging saurischian dinosaurs (Nesbitt, 2011; Stefanic and Nesbitt, 2019) and absent in the ornithischian *Scutellosaurus lawleri* found in the Kayenta Formation (Colbert, 1981; Breeden and Rowe, 2020; Breeden et al., 2021). If our identifications of the proximal ends of metatarsal III and metatarsal IV of ARCH 4012 are correct, the proximal shapes more closely resemble those of the neotheropod *Dilophosaurus wetherilli* than those of the sauropodomorph *Sarhsaurus aurifontanalis* (Figure 3), which are the two saurischians present in the Kayenta Formation in the same size range as ARCH 4012, suggesting a neotheropod affinity rather than that of a sauropodomorph.

Locality, Horizon, and Age

Locality ARCH 71v, near the Garden of Eden, about 2.5 km northwest of the Windows area of ARCH (Figure 1B); typical facies of the Kayenta Formation; “bone was found eroding from red mudstone next to exposed area of white sandstone” (Madsen et al., 2012, p. 16–17). The Kayenta Formation is Early Jurassic in age (Figure 1A), likely Pliensbachian–Sinemurian, and the middle of the silty facies of the Kayenta Formation in northeastern Arizona may be as young as about 183 Ma (Padian, 1989; Marsh et al., 2014; Suarez et al., 2017; Marsh, 2018).

DESCRIPTION

Axial Skeleton

Presacral Vertebrae

The partial centrum preserves less than half of what we hypothesize to be the posterior articular facet in posterior view (Figure 2A). We identify it as belonging to the posterior cervical or trunk series because the angle formed by the posterior and ventral margins in lateral view is rather acute; this angle is larger in anterior cervical centra that tend to be more parallelogram in shape in *Dilophosaurus wetherilli* (e.g., specimen UCMP 37302)




Figure 2. Caption is on the following page.

Figure 2 (figure is on the previous page). Preserved fragments of ARCH 4012 compared with elements of *Dilophosaurus wetherilli*, UCMP 37302. ARCH 4012, centrum face (A, B) in ?posterior (A) and ?right lateral view (C); UCMP 37302, vertebra 15 (C, D) in right lateral (C) and dorsal view (B); ARCH 4012, trunk prezygapophysis (E, F) in medial (E) and dorsal view (F); UCMP 37302, vertebra 54 in left lateral view (G); ARCH 4012, partial caudal centrum (H through J) in ?posterior (H), ?left lateral (I), and dorsal view (J); UCMP 37302, left radius in lateral view (K); ARCH 4012, distal end of ?left radius (L–N) in ?lateral (L), distal (M), and ?medial view (N); UCMP 37302, distal end of right femur, reversed (O, Q) in lateral (O) and distal view (Q); ARCH 4012, distal end of left femur (P, R) in lateral (P) and distal view (R); UCMP 37302, proximal end of left fibula (S, U) in proximal (S) and lateral view (U); ARCH 4012, fragment of ?left fibula shaft (T, V) in proximal (T) and lateral (V) view; UCMP 37302, proximal end of left metatarsal IV (W, Y) in lateral (W) and proximal view (Y); ARCH 4012, proximal end of ?left metatarsal IV (X, Z) in lateral (X) and proximal view (Z); UCMP 37302, proximal end of left metatarsal III (AA, AC) in ventral (AA) and proximal view (AC); ARCH 4012, proximal end of ?left metatarsal III (AB, AD) in ventral (AB) and proximal view (AD); UCMP 37302, right metatarsal I in lateral view (AE); ARCH 4012, distal end of ?right metatarsal I (AF, AG) in lateral (AF) and dorsal view (AG). Abbreviations: as, articular surface; hy, hypantrum; lc, lateral conyle; lf, ligament fossa; g, groove; ncj, neurocentral junction; prz, prezygapophysis; tfc, tibiofibular crest. Elements of UCMP 37302 are not to scale. Dashed lines indicate broken margins.




Figure 3. Left metatarsi of larger-bodied saurischians in the Kayenta Formation in proximal view. (A) *Dilophosaurus wetherilli*, UCMP 37302 (modified from Marsh and Rowe, 2020); (B) Theropoda, ARCH 4012; (C) *Sarahsaurus aurifontanalis*, TMM 43646-3 (reversed, modified from Marsh and Rowe, 2018). Elements are not to scale. Roman numerals identify each digit.

and *Sarahsaurus aurifontanalis* (e.g., specimen TMM 43646-2). The prezygapophysis preserves the relatively flat dorsal articular surface for the preceding postzygapophysis and the concave medial margin that would have formed the hypantrum (Figure 2F), which is present on the trunk vertebrae in *Dilophosaurus wetherilli* (e.g., UCMP 37302) and *Sarahsaurus aurifontanalis* (e.g., TMM 43646-2).

Caudal Vertebra

Less than half of the element is preserved as seen in lateral view (Figure 2I). The articular face is concave rather than flat, and the dorsal ridges that form the junction with the neural arch are present. The ventral margin of the articular face lacks the sloped surface that would have articulated with a haemal arch, indicating that this caudal vertebra was likely far enough back in the tail to be posterior to the last haemal arch.

Forelimb and Radius

The distal end of the ?left radius is identical to that of *Dilophosaurus wetherilli* (e.g., UCMP 77270) and *Sarahsaurus aurifontanalis* (e.g., TMM 43646-2). In lateral/medial view (Figures 2L and 2N), the posterior margin is relatively straight, and the anterior margin is concave. The distal outline is subelliptical (Figure 2M).

Hindlimb

Femur

Only the lateral sides of the lateral condyle and tibiofibular crests are preserved. A concavity is present on the lateral side of the tibiofibular crest that continued distally to form a groove between the tibiofibular crest and the lateral condyle (Figure 2P). In distal view the lateral margin of the lateral condyle is subcircular (Figure 2R).

Fibula

A portion of the proximal end of the shaft is present, identified by the cross-sectional shape in proximal view (Figure 2T). This is likely the shaft of a fibula and not a fragment of the shaft of the femur because what would be the medial side has a small patch of cortical bone that would not be present inside a femur. The fibula fragment would have been too proximal to preserve the ridge present in *Dilophosaurus wetherilli* (e.g., UCMP 37302) and *Sarhsaurus aurifontanalis* (e.g., TMM 43646-2) that represents the insertion site for the *M. iliofibularis* (Parrish, 1993; Nesbitt, 2011).

Metapodials

Half of the distal articular surface of metatarsal I is present (Figures 2AF and 2AG). It preserves a ligament fossa as well as the ventral expansion found in *Dilophosaurus wetherilli* (e.g., UCMP 37302) but not *Sarhsaurus aurifontanalis* (e.g., TMM 43646-3). The identifications of metatarsals III and IV are based on the shape of the partial proximal ends of each element (Figure 3B). Metatarsal III is broken ventrally but the dorsal margin is roughly parallelogram-shaped in proximal view (Figure 2AD), similar to that of *Dilophosaurus wetherilli* (e.g., UCMP 37302; Figure 3A). The shape of the proximal end is also consistent with that of *D. wetherilli* in ventral view. The dorsal and medial margins of the proximal end of metatarsal IV are complete, but the ventral portion is broken. In proximal view (Figure 2Z), the remaining bone is roughly subtriangular in shape, consistent with metatarsal IV of *Dilophosaurus wetherilli* (e.g., UCMP 37302).

DISCUSSION

Vertebrate Body Fossils and Age of the Kayenta Formation

Two major lithologic regimes are represented in the Kayenta Formation, described as the sandier ‘typical facies’ (i.e., what is present at the type section; Baker et al., 1936) and the finer-grained ‘silty facies’ (Averitt et al., 1955; Harshbarger et al., 1957). The delineation of these regimes bisects northern Arizona and southern

Utah (Figure 1B) such that the typical facies is present northeast of Tuba City, Arizona, including southeastern Utah, and the silty facies occurs at the Adeii Eechi Cliffs southeast of Tuba City and in southwestern Utah. These two facies have a dramatic difference in vertebrate body fossil occurrence, which drove differing age interpretations for the formation and the location of the Triassic-Jurassic boundary within the Glen Canyon Group since the early part of the 20th century.

Arizona

The Kayenta Formation was recognized as a different unit from the Todilto Formation of the Middle Jurassic San Rafael Group and ultimately named for its type section at Comb Ridge near Kayenta, Arizona (Baker et al., 1936). At that time, the formation was thought to be Triassic in age and its only reported fossils were plant “stems” (Baker et al., 1936, p. 5), unionid bivalves (Baker, 1933), and dinosaur footprints (*Eubrontes* and *Anchisauripus* according to R. Lull, Yale Peabody Museum, written communication, 1917; Gregory, 1917, p. 56). Age interpretations for the Kayenta Formation oscillated between the Late Triassic and Early Jurassic in subsequent decades, justified primarily by two major fossil discoveries within the formation published in the same year of 1954.

The first was the discovery of two skeletons of a relatively large hypothesized *Megalosaurus*-like theropod dinosaur from near the base of the silty facies near Moenave, Arizona (Welles, 1954); this taxon was later named *Dilophosaurus wetherilli* (Welles, 1970). Welles (1984) used the size and proportions of limb elements to argue Early Jurassic affinities for this new dinosaur, now hypothesized to be only distantly related to European Middle Jurassic tetanuran theropods (Marsh and Rowe, 2020). The type locality of *Dilophosaurus wetherilli* is across the road from a famous megatracksite at the base of the Kayenta Formation (Welles, 1971; Morales and Colbert, 1986; Lucas et al., 2005; Marsh, 2018).

The second major discovery was of a tritylodontid cynodont skeleton associated with a protosuchid crocodyliform (Lewis, 1958) near the top of the typical facies near Comb Ridge in Kayenta, Arizona (Figure 1B; Averitt et al., 1955; Harshbarger et al., 1957). The trity-

lodontid represented the first of its kind in the Western Hemisphere at the time and was first thought to more closely resemble 'Old World' taxa from the Triassic of China and South Africa (Averitt et al., 1955). Thus, during the late 1950s the silty facies of the Kayenta Formation seemed to provide a younger age estimate for the formation than the typical facies. Subsequent work helped to clarify this chronological uncertainty, including extensive revisions of the stratigraphic relationships and major unconformities of the Kayenta Formation and stratigraphically adjacent units (e.g., Lewis et al., 1961; Pippingos and O'Sullivan, 1978; Peterson and Pippingos, 1979; Clark and Fastovsky, 1986), more refined understanding of Triassic/Jurassic strata and vertebrate assemblages around the world (e.g., Olsen and Sues, 1986; Luo and Wu, 1994; Fang et al., 2000; Huang, et al., 2005; Bordy et al., 2020), revisions in the anatomy and taxonomy of theropods and tritylodontids (e.g., Kermack, 1982; Welles, 1984; Sues, 1986a, 1986b, 1986c), and the eventual use of U-Pb detrital zircon geochronology within the Glen Canyon Group (Dickinson and Gehrels, 2003, 2010; Marsh et al., 2014). The entire Kayenta Formation is now accepted to be Early Jurassic in age (Sinemurian-Pliensbachian) and the end-Triassic mass extinction and Triassic-Jurassic boundary (now chronologically decoupled; Blackburn et al., 2013) occur within the underlying Dinosaur Canyon Member of the Moenave Formation and the lateral equivalent of the Wingate Sandstone (Figure 1A; Lucas et al., 1997; Donohoo-Hurley et al., 2010; Martz et al., 2017; Suarez et al., 2017). Other than the tritylodontid (and others later found at the same site) and protosuchid mentioned above (Lewis, 1958), the only other vertebrate fossils from the typical facies of Arizona occur near Tsegi Canyon and include hybodont shark teeth and a potential batoid tooth (Curtis and Padian, 1999); however, it has been suggested that the latter is likely the result of screen wash contamination from a concurrent Cretaceous microvertebrate sampling effort at MNA (Milner et al., 2006).

The tritylodontid *Kayentatherium wellsi* was named from the silty facies near Many Farms, Arizona (Figure 1B; Kermack, 1982), and is considered the senior synonym of the only named tritylodontid from the typical facies of far northeastern Arizona (Lewis, 1956). Most

of the vertebrate assemblage of the Kayenta Formation is known from sites in the silty facies spanning the Echo and Adeii Eechii Cliffs along Ward Terrace (Figure 1B). This area includes the type localities of *Dilophosaurus wetherilli* (Welles, 1954), 'Syntarsus' *kayentakatae* (Rowe, 1989), *Sarhsaurus aurifontanalis* (Rowe et al., 2010), *Scutellosaurus lawleri* (Colbert, 1981), *Rhamphinion jenkinsi* (Padian, 1984), *Kayentasuchus walkeri* (Clark and Sues, 2002), *Eopneumatosuchus colberti* (Crompton and Smith, 1980), *Calsoyasuchus vallicept* (Tykoski et al., 2002), *Kayentachelys aprix* (Gaffney et al., 1987), *Navajosphenodon sani* (Simões et al., 2022), *Prosalirus bitis* (Shubin and Jenkins, 1995), *Eocaecilia macropodia* (Jenkins and Walsh, 1993), *Dinnebitodon amarali* (Sues, 1986b), and *Dinnetherium nezorom* (Jenkins et al., 1983). In addition to these taxa, the silty facies in that area also produced a larger-bodied thyreophoran (Padian, 1989), unnamed protosuchid crocodyliforms (Clark and Sues, 2002), an *Oligokyphus*-like tritylodontid (Sues, 1985), a lungfish (Kirkland, 1987; Milner and Kirkland, 2006), and microvertebrates including actinopterygians (Clark and Fastovsky, 1986; Curtis and Padian, 1999; Milner et al., 2006), hybodontoid sharks, potential salamanders, and a haramiyid mammaliaform (Curtis and Padian, 1999).

Utah

The Kayenta Formation in Utah includes the typical facies in the eastern region near Moab and the silty facies in the western region around St. George (Figure 1B; Harshbarger et al., 1957; Priddy and Clarke, 2020), the latter being underlain by the Springdale Sandstone as a second lower unit in the formation (Lucas and Tanner, 2006). The Springdale Sandstone, the silty facies, and the typical facies of the Kayenta Formation frequently preserve trace fossils of various dinosaurs and other archosaurs, including megatracksites in the St. George area (DeBlieux et al., 2004, 2006; Hamblin et al., 2006; Milner et al., 2012 and references therein; Lockley et al., 2018).

The silty facies of Utah preserves semionotid fish scales at Zion National Park (Lockley, 1984; DeBlieux et al., 2004, 2006; Milner and Kirkland, 2006), coelacanths, semionotid and paleoniscoid fish, and poten-

tial crocodylomorph bones at Warner Valley (Milner et al., 2012), as well as unpublished articulated semionotids and teeth belonging to actinopterygians, crocodylomorphs, ornithischians, theropods, and a possible pterosaur from a more recent roadcut near St. George (Figure 1B; Milner et al., 2017). The typical facies also preserves other fish in Utah, including a semionotid fish skeleton from near Moab (Milner et al., 2006) on exhibit at the Moab Museum, and a lungfish tooth plate west of Goblin Valley (Frederickson and Cifelli, 2017). Despite occurring within what would otherwise be mapped as the typical facies of the Kayenta Formation (Figure 1B), the ARCH dinosaur described here occurs in interbedded red mudstone and white sandstone in the northern part of the Paradox Basin (Madsen et al., 2012), which may have been providing more syndepositional accommodation owing to salt dissolution in the underlying Pennsylvanian Paradox Formation and thus deposit more silt than would otherwise be expected in the typical facies in the broader region (Bromley, 1991).

Other Dinosaur Fossils from Arches National Park

Dinosaur fossils are relatively common in the Mesozoic units within the boundaries of ARCH and are mostly represented by tracks. The Church Rock Member of the Upper Triassic Chinle Formation preserves the ichnogenus *Grallator*, likely made by a small-bodied coelophysoid theropod (Lockley and Gierliński, 2009; Martz et al., 2017). *Grallator* is also known from the base of the overlying Wingate Sandstone (Swanson et al., 2005; Madsen et al., 2012). At least the lower part of the Wingate Sandstone is Late Triassic in age owing to the presence of a phytosaur in the nearby Lisbon Valley south of Moab (Morales and Ash, 1993; Martz et al., 2017). A single zygapophysis was found as float during a survey of the Wingate Sandstone (Madsen et al., 2012), but it is rather large for contemporaneous dinosaurs and because the source stratum was not found it is more parsimonious to consider it to have come down from the overlying Kayenta given the rarity of body fossils from the Wingate Sandstone (Tweet and Santucci, 2018).

In addition to the ARCH dinosaur body fossils

described here, the Kayenta Formation also preserves indeterminate theropod tracks (Swanson et al., 2005), and the overlying Lower Jurassic Navajo Sandstone includes *Grallator* (Madsen et al., 2012) and tracks of sauropodomorph dinosaurs (*Navahopus* or *Otozoum*; Tweet et al., 2018). In the lower Upper Jurassic units there are tridactyl tracks within the Moab Tongue Member of the Curtis Formation (Swanson et al., 2005). The famous Moab Megatracksite (including ARCH) at the contact of the Moab Tongue Member and the overlying Summerville Formation preserves innumerable tracks and trackways referred to the larger ichnogenera *Megalosauripus* and *Therangospodus* (Lockley, 1991, 2022). Higher in the section, the Upper Jurassic Morrison Formation preserves sauropod postcranial bones including a partial apatosaurine skeleton (Doelling, 1985; Swanson, et al. 2005). The Ruby Ranch Member of the Lower Cretaceous Cedar Mountain Formation preserves numerous dinosaurian ichnotaxa, including those belonging to iguanodontid ornithischians, sauropods, and theropods including dromaeosaurids (Swanson et al., 2005; Madsen et al., 2012; Lockley et al., 2014).

CONCLUSIONS

The ARCH specimen described here is important because (1) it represents the first body fossils of a dinosaur from the Kayenta Formation of Utah outside of teeth from a roadcut locality near St. George, and (2) it represents the only unambiguous dinosaur body fossil from the Kayenta Formation within Arches National Park. The U.S. National Park Service has 286 units known to contain fossils (67% of all units as of March 2023; <https://www.nps.gov/subjects/fossils/fossil-parks-list.htm>) and not all have been surveyed, but only 16 were established explicitly to protect paleontological resources (Tweet and Santucci, 2021). ARCH was founded to protect geologic resources, and though paleontological resources are considered important resources, they are not considered fundamental resources to the park (https://www.nps.gov/arch/learn/management/foundation-document.htm#CP_JUMP_5740030). This discovery shows the potential for more vertebrate body fossils from the typical facies of the Kayenta in Arizona and Utah, highlighting the importance of inventory

and monitoring of fossil resources in federally protected areas.

ACKNOWLEDGMENTS

Thanks to Scott Madsen (UGS) and Vince Santucci (NPS) for their work in the field and reporting on the 2011 ARCH paleontological survey. We thank Kevin Padian, Pat Holroyd, and Mark Goodwin (UCMP), Chris Sagebiel and Matt Brown (TMM), Peekay Briggs (ARCH), and Matt Smith and Debbie Wagner (PEFO) for access to collections and archives in their care. This is Petrified Forest National Park Paleontological Contribution No. 90. The conclusions presented here are those of the authors and do not represent the views of the United States Government. Funding was provided by the Utah Geological Survey and the National Park Service. We thank the reviewers and editors of this article for their helpful comments and suggestions.

REFERENCES

- Averitt, P., Detterman, J.S., Harshbarger, J.W., Repenning, C.A., and Wilson, R.F., 1955, Revisions in correlation and nomenclature of Triassic and Jurassic formations in southwestern Utah and northern Arizona: *Bulletin of the American Association of Petroleum Geologists*, v. 39, p. 2515–2535, doi: 10.1306/5CEAE2E9-16BB-11D7-8645000102C1865D.
- Baker, A.A., 1933, Geology and oil possibilities of the Moab district, Grand and San Juan Counties, Utah: *U.S. Geological Survey Bulletin*, v. 841, p. 1–95, doi: 10.3133/b841.
- Baker, A.A., Dane, C.H., and Reeside, J.B., 1936, Correlation of the Jurassic formations of parts of Utah, Arizona, New Mexico and Colorado: *U.S. Geological Survey Professional Paper*, v. 183, p. 1–66, doi: 10.3133/pp183.
- Benton, M.J., 1999, *Scleromochlus taylori* and the origin of dinosaurs and pterosaurs: *Philosophical Transactions of the Royal Society of London*, v. 354, p. 1423–1446, doi: 10.1098/rstb.1999.0489.
- Blackburn, T.J., Olsen, P.E., Bowring, S.A., McLean, N.M., Kent, D.V., Puffer, J., McHone, G., Rasbury, E.T., and Et-Touami, M., 2013, Zircon U-Pb geochronology links the end-Triassic extinction with the Central Atlantic Magmatic Province: *Science*, v. 340, p. 941–945, doi: 10.1126/science.1234204.
- Blakey, R.C., 1994, Paleogeographic and tectonic controls on some Lower and Middle Jurassic erg deposits, Colorado Plateau, in Caputo, M.V., Peterson, J.A., and Franczyk, K.J., editors, *Mesozoic systems of the Rocky Mountain region*, USA: Society for Sedimentary Geology (SEPM), Rocky Mountain Section Publication, p. 273–298.
- Bordy, E.M., Abrahams, M., Sharman, G.R., Viglietti, P.A., Benson, R.B.J., McPhee, B.W., Barrett, P.M., Sciscio, L., Condon, D., Mundil, R., Raderman, Z., Jinnah, Z., Clark, J.M., Suarez, C.A., Chapelle, K.E.J., and Choiniere, J.N., 2020, A chronostratigraphic framework for the upper Stormberg Group—implications for the Triassic-Jurassic boundary in southern Africa: *Earth-Science Reviews*, v. 203, p. 103120, doi: 10.1016/j.earscirev.2020.103120.
- Breeden III, B.T., and Rowe, T.B., 2020, Specimens of *Scutellosaurus lawleri* Colbert, 1981, from the Lower Jurassic Kayenta Formation in Arizona elucidate the early evolution of thyreophoran dinosaurs: *Journal of Vertebrate Paleontology*, v. 40, p. e1791894, doi: 10.1080/02724634.2020.1791894.
- Breeden III, B.T., Raven, T.J., Butler, R.J., Rowe, T.B., and Maidment, S.C.R., 2021, The anatomy and palaeobiology of the early armoured dinosaurs *Scutellosaurus lawleri* (Ornithischia: Thyreophora) from the Kayenta Formation (Lower Jurassic) of Arizona: *Royal Society Open Science*, v. 8, p. 201676, doi: 10.1098/rsos.201676.
- Bromley, M.H., 1991, Architectural features of the Kayenta Formation (Lower Jurassic), Colorado Plateau, USA—relationship to salt tectonics in the Paradox Basin: *Sedimentary Geology*, v. 74, p. 77–99, doi: 10.1016/0037-0738(91)90024-8.
- Clark, J.M., and Fastovsky, D.E., 1986, Vertebrate biostratigraphy of the Glen Canyon Group in northern Arizona, in Padian, K., editor, *The beginning of the Age of Dinosaurs—faunal change across the Triassic-Jurassic boundary*: Cambridge University Press, p. 285–301.
- Clark, J.M., and Sues, H.-D., 2002, Two new basal crocodylomorph archosaurs from the Lower Jurassic and the monophyly of the Sphenosuchia: *Zoological Journal of the Linnean Society*, v. 136, p. 77–95, doi: 10.1046/j.1096-3642.2002.00026.x.
- Colbert, E.H., 1981, A primitive ornithischian dinosaur from the Kayenta Formation of Arizona: *Museum of Northern Arizona Bulletin*, v. 53, p. 1–61.
- Crompton, A.W., and Smith, K.K., 1980, A new genus and species of crocodylian from the Kayenta Formation (Late Triassic?) of northern Arizona, in Jacobs, L.L., editor, *Aspects of vertebrate history—essays in honor of Edwin Harris Colbert*: Museum of Northern Arizona Press, p. 193–217.
- Curtis, K., and Padian, K., 1999, An Early Jurassic microvertebrate fauna from the Kayenta Formation of northeastern Arizona—microfaunal change across the Triassic-Jurassic boundary: *PaleoBios*, v. 19, p. 19–37.
- De Blieux, D.D., Kirkland, J.I., Smith, J.A., McGuire, J., and San-

- tucci, V.L., 2006, An overview of the paleontology of Upper Triassic and Lower Jurassic rocks in Zion National Park, Utah: New Mexico Museum of Natural History and Science Bulletin, v. 37, p. 490–501.
- De Blieux, D.D., Smith, J.A., McGuire, J.L., Kirkland, J.I., and Santucci, V.L., 2004, Zion National Park Paleontological Survey: Utah Geological Survey Contract Deliverable, 89 p.
- Dickinson, W.R., and Gehrels, G.E., 2003, U-Pb ages of detrital zircons from Permian and Jurassic eolian sandstones of the Colorado Plateau, USA—paleogeographic implications: *Sedimentary Geology*, v. 163, p. 29–66, doi: 10.1016/S0037-0738(03)00158-1.
- Dickinson, W.R., and Gehrels, G.E., 2010, Insights into North American paleogeography and paleotectonics from U-Pb ages of detrital zircons in Mesozoic strata of the Colorado Plateau, USA: *International Journal of Earth Sciences*, v. 99, p. 1247–1265, doi: 10.1007/s00531-009-0462-0.
- Doelling, H.H., 1985, Geology of Arches National Park: Utah Geological and Mineral Survey Map 74, 15 p. pamphlet, scale 1:50,000.
- Donohoo-Hurley, L.L., Geissman, J.W., and Lucas, S.G., 2010, Magnetostratigraphy of the uppermost Triassic and lowermost Jurassic Moenave Formation, Western United States—correlation with strata in the United Kingdom, Morocco, Turkey, Italy, and Eastern United States: *Geological Society of America Bulletin*, v. 122, p. 2005–2019, doi: 10.1130/B30136.1.
- Fang, X.S., Pang, Q.J., Lu, L.W., Zhang, Z.X., Pan, S.G., Wang, Y.M., Li, X.K., and Cheng, Z.W., 2000, Lower, Middle, and Upper Jurassic subdivision in the Lufeng region, Yunnan Province, in *Proceedings of the Third National Stratigraphical Congress of China*: Beijing, China, Geological Publishing House, p. 208–214.
- Frederickson, J.A., and Cifelli, R.L., 2017, New cretaceous lungfishes (Dipnoi, Ceratodontidae) from Western North America: *Journal of Paleontology*, v. 91, p. 146–161, doi: 10.1017/jpa.2016.131.
- Gaffney, E.S., Hutchison, H., Jenkins, Jr., F.A., and Meeker, L.J., 1987, Modern turtle origins—the oldest known cryptodire: *Science*, v. 237, p. 289–291, doi: 10.1126/science.237.4812.289.
- Gauthier, J.A., 1986, Saurischian monophyly and the origin of birds: *Memoirs of the California Academy of Sciences*, v. 8, p. 1–55.
- Gregory, H.E., 1917, Geology of the Navajo country—a reconnaissance of parts of Arizona, New Mexico, and Utah: U.S. Geological Survey Professional Paper, v. 93, p. 1–161, doi: 10.3133/pp93.
- Hamblin, A.H., Lockley, M.G., and Milner, A.R.C., 2006, More reports of theropod dinosaur tracksites from the Kayenta Formation (Lower Jurassic), Washington County, Utah—implications for describing the Springdale Megatracksite: *New Mexico Museum of Natural History and Science Bulletin*, v. 37, p. 276–281.
- Harshbarger, J.W., Repenning, C.A., and Irwin, J.H., 1957, Stratigraphy of the uppermost Triassic and the Jurassic rocks of the Navajo Country: U.S. Geological Survey Professional Paper, v. 291, p. 1–74, doi: 10.3133/pp291.
- Huang, B.C., Li, Y.A., Fang, X.S., Sun, D.J., Pang, Q.Q., Cheng, Z.W., and Li, P.W., 2005, Magnetostratigraphy of the Jurassic in Lufeng, central Yunnan: *Geological Bulletin of China*, v. 24, p. 322–328.
- Hunt, A.P., and Lucas, S.G., 2006, Triassic-Jurassic tetrapod ichnofacies: *New Mexico Museum of Natural History and Science Bulletin*, v. 37, p. 12–22.
- Jenkins, F.A., Jr., and Walsh, D.M., 1993, An Early Jurassic caecilian with limbs: *Nature*, v. 365, p. 246–250, doi: 10.1038/365246a0.
- Jenkins, F.A., Jr., Crompton, A.W., and Downs, W.R., 1983, Mesozoic mammals from Arizona—new evidence on mammalian evolution: *Science*, v. 222, p. 1233–1235, doi: 10.1126/science.222.4629.12.
- Kermack, D.M., 1982, A new tritylodontid from the Kayenta Formation of Arizona: *Zoological Journal of the Linnean Society*, v. 76, p. 1–17, doi: 10.1111/j.1096-3642.1982.tb01953.x.
- Kirkland, J.I., 1987, Upper Jurassic and Cretaceous lungfish tooth plates from the Western Interior, the last dipnoan faunas of North America: *Hunteria*, v. 2, p. 1–26.
- Lewis, G.E., 1956, *Nearctylon broomi*, the first Nearctic tritylodontid, in Hotton, N., III, MacLean, P.D., Roth, J.J., and Roth, E.C., editors, *The ecology and biology of mammal-like reptiles*: Smithsonian Institution Press, p. 295–303.
- Lewis, G.E., 1958, American Triassic mammal-like vertebrates: *Geological Society of America Bulletin*, v. 69, p. 1735.
- Lewis, G.E., Irwin, J.H., and Wilson, R.F., 1961, Age of the Glen Canyon Group (Triassic and Jurassic) on the Colorado Plateau: *Geological Society of America Bulletin*, v. 71, p. 1437–1440, doi: 10.1130/0016-7606(1961)72[1437:AOTG-CG]2.0.CO;2.
- Lockley, M.G., 1984, Dinosaur footprints from the Kayenta Formation of eastern Utah: *University of Colorado at Denver, Geology Department Magazine*, Fall 1984, p. 29–32.

- Lockley, M.G., 1991, The Moab Megatracksite—a preliminary description and discussion of millions of Middle Jurassic tracks in eastern Utah, in Averett, W.R., editor, Guidebook for dinosaur quarries and tracksites tour, western Colorado and eastern Utah: Grand Junction Geological Society, p. 59–65.
- Lockley, M.G., 2022, The distribution of theropod-dominated ichnofaunas in the Moab Megatracksite area, Utah: implications for Late Jurassic palaeoecology along an arid coast: *Historical Biology*, v. 34, p. 1717–1751 (doi: 10.1080/08912963.2021.1975279).
- Lockley, M.G., and Gierliński, G.D., 2009, A *Grallator*-dominated tracksite from the Chinle Group (Late Triassic), Moab, Utah: *Geological Quarterly*, v. 53, p. 433–440.
- Lockley, M.G., Gierliński, G.D., Dubicka, Z., Breithaupt, B.H., and Matthews, N.A., 2014, A preliminary report on a new dinosaur tracksite in the Cedar Mountain Formation (Cretaceous) of eastern Utah: *New Mexico Museum of Natural History and Science Bulletin*, v. 62, p. 279–286.
- Lockley, M.G., and Hunt, A.P., 1994, A review of Mesozoic vertebrate ichnofaunas of the Western Interior United States—evidence and implications of a superior track record, in Caputo, M.V., Peterson, J.A., and Franczyk, J.F., editors, Mesozoic systems of the Rocky Mountain region, USA: Society for Sedimentary Geology (SEPM), Rocky Mountain Section Publication, p. 95–108.
- Lockley, M.G., and Hunt, A.P., 1995, Dinosaur tracks and other fossil footprints of the Western United States: New York, Columbia University Press, 360 p.
- Lockley, M.G., Matthews, N.A., Breithaupt, B.H., Gierliński, G., Cart, K., and Hunt-Foster, R., 2018, Dinosaur tracksites in the Lower Jurassic Kayenta Formation near Moab, Utah—implications for paleoecology: *New Mexico Museum of Natural History and Science Bulletin*, v. 79, p. 441–449.
- Lucas, S.G., and Tanner, L.H., 2006, The Springdale Member of the Kayenta Formation, Lower Jurassic of Utah-Arizona: *New Mexico Museum of Natural History and Science Bulletin*, v. 37, p. 71–76.
- Lucas, S.G., Heckert, A.B., Anderson, O.J., and Estep, J.W., 1997, Phytosaur from the Wingate Sandstone in southeastern Utah and the Triassic-Jurassic boundary on the Colorado Plateau, in Mesa Southwest Museum and Southwest Paleontological Society, Proceedings: Mesa, Arizona, Mesa Southwestern Museum, v. 1, p. 49–59.
- Lucas, S.G., Heckert, A.B., and Tanner, L.H., 2005, Arizona's Jurassic vertebrates and the age of the Glen Canyon Group: *New Mexico Museum of Natural History and Science Bulletin*, v. 29, p. 95–104.
- Luo, Z., and Wu, X.-C., 1994, The small tetrapods of the lower Lufeng Formation, Yunnan, China, in N. C. Fraser and H.-D. Sues, editors, In the Shadow of the Dinosaurs: Early Mesozoic Tetrapods: Cambridge University Press, p. 251–270.
- Madsen, S.K., Kirkland, J.I., De Blieux, D.D., Santucci, V.L., Inkenbrandt, P., and Tweet, J.S., 2012, Paleontological resources inventory and monitoring, Arches National Park, Utah: Utah Geological Survey Contract Deliverable, Cooperative Agreement #H230097080, 119 p.
- Marsh, A.D., 2018, Contextualizing the evolution of theropod dinosaurs in Western North America using U-Pb geochronology of the Chinle Formation and Kayenta Formation on the Colorado Plateau: Austin, The University of Texas, M.S. thesis, 333 p.
- Marsh, A.D., and Rowe, T.B., 2018, Anatomy and systematics of the sauropodomorph *Sarawsaurus aurifontanalis* from the Early Jurassic Kayenta Formation: *PLoS ONE*, v. 13, p. e0204007, doi: 10.1371/journal.pone.0204007.
- Marsh, A.D., and Rowe, T.B., 2020, A comprehensive anatomical and phylogenetic evaluation of *Dilophosaurus wetherilli* (Dinosauria, Theropoda) with descriptions of new specimens from the Kayenta Formation of northern Arizona: *Journal of Paleontology*, v. 94 (Supplement 78), p. 1–103, doi: 10.1371/journal.pone.0204007.
- Marsh, A.D., Rowe, T., Simonetti, A., Stockli, D., and Stockli, L., 2014, The age of the Kayenta Formation of northeastern Arizona—overcoming the challenges of dating fossil bone [abs.]: *Journal of Vertebrate Paleontology, Program and Abstracts*, p. 178.
- Marsh, O.C., 1881, Principal characters of American Jurassic dinosaurs, part V: *American Journal of Science*, v. 21 (Series 3), p. 417–423, doi: 10.2475/ajs.s3-21.125.417.
- Martz, J.W., Kirkland, J.I., Milner, A.R.C., Parker, W.G., and Santucci, V.L., 2017, Upper Triassic lithostratigraphy, depositional systems, and vertebrate paleontology across southern Utah: *Geology of the Intermountain West*, v. 4, p. 99–180, doi: 10.31711/giw.v4.pp99-180.
- Milner, A.R.C., and Kirkland, J.I., 2006, Preliminary review of the Early Jurassic (Hettangian) freshwater Lake Dixie fish fauna in the Whitmore Point Member, Moenave Formation in southwest Utah: *New Mexico Museum of Natural History and Science Bulletin*, v. 37, p. 510–521.
- Milner, A.R.C., Kirkland, J.I., and Borthisell, T.A., 2006, The geographic distribution and biostratigraphy of Late Triassic-Early Jurassic freshwater fish faunas of the southwestern United States: *New Mexico Museum of Natural History and Science Bulletin*, v. 37, p. 522–529.

- Milner, A.R.C., Birthisel, T.A., Kirkland, J.I., Breithaupt, B.H., Matthews, N.A., Lockley, M.G., Santucci, V.L., Gibson, S.Z., De Blieux, D.D., Hurlbut, M., Harris, J.D., and Olsen, P.E., 2012, Tracking Early Jurassic dinosaurs across southwestern Utah and the Triassic-Jurassic transition, in Bonde, J.W., and Milner, A.R.C., editors, Nevada State Museum paleontological papers: Nevada Department of Tourism and Cultural Affairs, v. 1, p. 1–107.
- Milner, A.R.C., Gay, R.J., Irmis, R.B., Overkamp, F., and Santella, M., 2017, New southwestern Utah paleontological locality from the Lower Jurassic Kayenta Formation reveals a diverse vertebrate fauna based on teeth and tracks [abs.]: Journal of Vertebrate Paleontology, Program and Abstracts, p. 164.
- Morales, M., and Ash, S.R., 1993, The last phytosaurs?: New Mexico Museum of Natural History and Science Bulletin, v. 3, p. 357–358.
- Morales, M., and Colbert, E.H., 1986, Stratigraphic occurrences of dinosaur tracks in the Glen Canyon Group of Arizona [abs.], in Gillette, D.D., editor, First International Symposium of dinosaur tracks and traces, abstracts with program, New Mexico, Albuquerque: New Mexico Museum of Natural History and Science, p. 21.
- National Park Service, Arches National Park, undated, Foundation document: online, https://www.nps.gov/arch/learn/management/foundation-document.htm#CP_JUMP_5740030, accessed December 20, 2023.
- National Park Service, Geological Resources Division, 2023, Fossil parks master list: online, <https://www.nps.gov/subjects/fossils/fossil-parks-list.htm>, accessed December 20, 2023.
- Nesbitt, S.J., 2011, Early evolution of archosaurs: relationships and the origin of major clades: Bulletin of the American Museum of Natural History, v. 352, p. 1–292, doi: 10.1206/352.1.
- Nesbitt, S.J., Butler, R.J., Ezcurra, M.D., Barrett, P.M., Stocker, M.R., Angielczyk, K.D., Smith, R.M., Sidor, C.A., Niedźwiedzki, G., Sennikov, A.G., and Charig, A.J., 2017, The earliest bird-line archosaurs and the assembly of the dinosaur body plan: Nature, v. 544, p. 484–487, doi: 10.1038/nature22037.
- Olsen, P.E., and Sues, H.-D., 1986, Correlation of continental Late Triassic and Early Jurassic sediments, and patterns of the Triassic-Jurassic tetrapod transition, in Padian, K., editor, The beginning of the Age of Dinosaurs—faunal change across the Triassic-Jurassic boundary: Cambridge University Press, p. 321–351.
- Owen, R., 1842, Report on British fossil reptiles, part II: Report of the British Association of the Advancement of Science, v. 11, p. 60–104.
- Padian, K., 1984, Pterosaur remains from the Kayenta Formation (?Early Jurassic) of Arizona: Palaeontology, v. 27, p. 407–413.
- Padian, K., 1989, Presence of the dinosaur *Scelidosaurus* indicates Jurassic age for the Kayenta Formation (Glen Canyon Group, northern Arizona): Geology, v. 17, p. 438–441, doi: 10.1130/0091-7613(1989)017<0438:POTDSI>2.3.CO;2.
- Padian, K., and May, C.L., 1993, The earliest dinosaurs: New Mexico Museum of Natural History and Science Bulletin, v. 3, p. 379–381.
- Parrish, J.M., 1993, Phylogeny of the Crocodylotarsi, with reference to archosaurian and crurotarsan monophyly: Journal of Vertebrate Paleontology, v. 13, p. 287–308, doi: 10.1080/02724634.1993.10011511.
- Peterson, F., and Pippingos, G.N., 1979, Stratigraphic relations of the Navajo Sandstone to Middle Jurassic Formations, southern Utah and northern Arizona: U.S. Geological Survey Professional Paper, v. 1035-B, p. 1–43, doi: 10.3133/pp1035B.
- Pippingos, G.N., and O'Sullivan, R.B., 1978, Principal unconformities in Triassic and Jurassic rocks, Western Interior United States—a preliminary survey: U.S. Geological Survey Professional Paper, v. 1035-A, p. 1–29, doi: 10.3133/pp1035A.
- Priddy, C.L. and Clarke, S.M., 2020, The sedimentology of an ephemeral fluvial-aeolian succession: Sedimentology, v. 67, p. 2392–2425, doi: 10.1111/sed.12706.
- Riggs, N.R., and Blakey, R.C., 1993, Early and Middle Jurassic paleogeography and volcanology of Arizona and adjacent areas, in Dunne G.C., and McDougall, K.A., editors, Mesozoic paleogeography of the Western United States II: Society for Sedimentary Geology (SEPM), p. 347–375.
- Riggs, N.R., Mattinson, J.M., and Busby, C.J., 1993, Correlation of Jurassic eolian strata between the magmatic arc and the Colorado Plateau—new U-Pb geochronology data from southern Arizona: Geological Society of America Bulletin, v. 105, p. 1231–1246, doi: 10.1130/0016-7606(1993)105<1231:COJESB>2.3.CO;2.
- Rowe, T.B., 1989, A new species of the theropod dinosaur *Syntarsus* from the Early Jurassic Kayenta Formation of Arizona: Journal of Vertebrate Paleontology, v. 9, p. 125–136, doi: 10.1080/02724634.1989.10011748.
- Rowe, T.B., Sues, H.-D., and Reisz, R.R., 2010, Dispersal and diversity in the earliest North American sauropodomorph dinosaurs, with a description of a new taxon: Proceedings

- of the Royal Society B, Biological Sciences, v. 278, p. 1044–1053, doi: 10.1098/rspb.2010.1867.
- Seeley, H.G., 1887, On the classification of the fossil animals commonly named Dinosauria: Proceedings of the Royal Society of London, v. 43, p. 165–171.
- Shubin, N.H., and Jenkins, F.A., Jr., 1995, An Early Jurassic jumping frog: *Nature*, v. 377, p. 49–52, doi: 10.1038/377049a0.
- Simões, T.R., Kinney-Broderick, G., and Pierce, S.E., 2022, An exceptionally preserved *Sphenodon*-like sphenodontian reveals deep time and conservation of the tuatara skeleton and ontogeny: *Communications Biology*, v. 5, p. 1–19, doi: 10.1038/s42003-022-03144-y.
- Stefanic, C.M., and Nesbitt, S.J., 2019, The evolution and role of the hyposphene-hypantrum articulation in Archosauria—phylogeny, size and/or mechanics: *Royal Society Open Science*, v. 6, p. 190258, doi: 10.1098/rsos.190258.
- Suarez, C.A., Knobbe, T.K., Crowlet, J.L., Kirkland, J.I., and Milner, A.R.C., 2017, A chronostratigraphic assessment of the Moenave Formation, USA using C-isotope chemostratigraphy and detrital zircon geochronology—implications for the terrestrial end Triassic extinction: *Earth and Planetary Science Letters*, v. 475, p. 83–93, doi: 10.1016/j.epsl.2017.07.028.
- Sues, H.-D., 1985, First record of the tritylodontid *Oligokyphus* (Synapsida) from the Lower Jurassic of Western North America: *Journal of Vertebrate Paleontology*, v. 5, p. 328–335, doi: 10.1080/02724634.1985.10011869.
- Sues, H.-D., 1986a, The skull and dentition of two tritylodontid synapsids from the Lower Jurassic of Western North America: *Bulletin of the Museum of Comparative Zoology*, v. 151, p. 217–268.
- Sues, H.-D., 1986b, *Dinnebitodon amarali*, a new tritylodontid (Synapsida) from the Lower Jurassic of Western North America: *Journal of Paleontology*, v. 60, p. 758–762, doi: 10.1017/S0022336000022277.
- Sues, H.-D., 1986c, Relationships and biostratigraphic significance of the Tritylodontidae (Synapsida) from the Kayenta Formation of northeastern Arizona, in Padian, K., editor, *The beginning of the Age of Dinosaurs—faunal change across the Triassic-Jurassic boundary*: Cambridge University Press, p. 277–284.
- Sues, H.-D., Clark, J.M., and Jenkins, F.A., Jr., 1994, A review of the Early Jurassic tetrapods from the Glen Canyon Group of the American Southwest, in Fraser, N.C., and Sues, H.-D., editors, *In the shadow of the dinosaurs—early Mesozoic tetrapods*: Cambridge University Press, p. 284–294.
- Swanson, B.A., Santucci, V.L., Madsen, S.K., Elder, A.S., and Kenworthy, J.P., 2005, Arches National Park paleontological survey: National Park Service Technical Report, NPS/NRGRD/GRDTR-05/01.
- Tweet, J.S., and Santucci, V.L., 2018, An inventory of non-avian dinosaurs from National Park Service areas: *New Mexico Museum of Natural History and Science Bulletin*, v. 79, p. 703–730.
- Tweet, J.S., and Santucci, V.L., 2021, From microfossils to megafauna—an overview of the taxonomic diversity of National Park Service fossils: *New Mexico Museum of Natural History and Science Bulletin*, v. 82, p. 437–457.
- Tykoski, R.S., 2005, Vertebrate paleontology in the Arizona Jurassic: *Mesa Southwest Museum Bulletin*, v. 11, p. 72–93.
- Tykoski, R.S., Rowe, T.B., Ketcham, R.A., and Colbert, M.W., 2002, *Calsoyasuchus valliceps*, a new crocodyliform from the Early Jurassic Kayenta Formation of Arizona: *Journal of Vertebrate Paleontology*, v. 22, p. 593–611, doi: 10.1671/0272-4634(2002)022[0593:CVANCF]2.0.CO;2.
- Welles, S.P., 1954, New Jurassic dinosaur from the Kayenta Formation of Arizona: *Bulletin of the Geological Society of America*, v. 65, p. 591–598, doi: 10.1130/0016-7606(1954)65[591:NJDFTK]2.0.CO;2.
- Welles, S.P., 1970, *Dilophosaurus* (Reptilia: Saurischia), a new name for a dinosaur: *Journal of Paleontology*, v. 44, p. 989.
- Welles, S.P., 1971, Dinosaur footprints from the Kayenta Formation of northern Arizona: *Plateau*, v. 4, p. 27–38.
- Welles, S.P., 1984, *Dilophosaurus wetherilli* (Dinosauria, Theropoda), osteology and comparisons: *Palaeontographica Abteilung A*, v. 185, p. 85–180.